

FINITE ENTROPY CHARACTERIZES TOPOLOGICAL RIGIDITY ON CONNECTED GROUPS

SIDDHARTHA BHATTACHARYA AND THOMAS WARD

ABSTRACT. Let X_1, X_2 be mixing connected algebraic dynamical systems with the Descending Chain Condition. We show that every equivariant continuous map $X_1 \rightarrow X_2$ is affine (that is, X_2 is *topologically rigid*) if and only if the system X_2 has finite topological entropy.

1. INTRODUCTION

An *algebraic $\mathbb{Z}^d$ -action* α on a compact abelian group X is a homomorphism $\alpha : \mathbf{n} \mapsto \alpha(\mathbf{n})$ from $\mathbb{Z}^d$ to the group $\text{Aut}(X)$ of continuous automorphisms of X . Compact groups are assumed to be metrizable throughout and are written multiplicatively; e is used to denote the identity element of any group. Write $X = (X, \alpha)$ for such an algebraic dynamical system, and call the system X connected, mixing and so on if X is connected, α is mixing, and so on.

Any algebraic system X preserves λ_X , the Haar measure on X . The system X is *mixing* if

$$\lim_{\mathbf{n} \rightarrow \infty} \lambda_X(A_1 \cap \alpha(\mathbf{n})(A_2)) = \lambda_X(A_1) \cdot \lambda_X(A_2)$$

for all measurable sets $A_1, A_2 \subset X$.

A map $\phi : X_1 \rightarrow X_2$ between algebraic dynamical systems is *equivariant* if $\phi \circ \alpha_1(\mathbf{n}) = \alpha_2(\mathbf{n}) \circ \phi$ for all $\mathbf{n} \in \mathbb{Z}^d$, and is *affine* if there is a continuous group homomorphism $\psi : X_1 \rightarrow X_2$ and an element $y \in X_2$ with $\phi(x) = \psi(x) \cdot y$.

Topological (respectively, measurable) rigidity is a property of the target system X_2 that forces an equivariant continuous (resp. measurable) map to coincide everywhere (resp. almost everywhere) with an affine map.

For $d \geq 1$, denote by $R_d = \mathbb{Z}[u_1^{\pm 1}, \dots, u_d^{\pm 1}]$ the ring of Laurent polynomials with integral coefficients in d commuting variables $u_1, \dots, u_d$. An element f of R_d is written

$$f(\mathbf{u}) = \sum_{\mathbf{n} \in \mathbb{Z}^d} f_{\mathbf{n}} \mathbf{u}^{\mathbf{n}}$$

1991 *Mathematics Subject Classification.* 22D40, 37B40.

The first author gratefully acknowledges the support of London Mathematical Society grant 4714 and the hospitality of the University of East Anglia.

with $\mathbf{u}^{\mathbf{n}} = u_1^{n_1} \cdots u_d^{n_d}$, $f_{\mathbf{n}} \in \mathbb{Z}$ for all $\mathbf{n} = (n_1, \dots, n_d) \in \mathbb{Z}^d$, and $f_{\mathbf{n}} = 0$ for all but finitely many $\mathbf{n} \in \mathbb{Z}^d$.

If $\mathbf{X} = (X, \alpha)$ is an algebraic $\mathbb{Z}^d$ -action on a compact abelian group X , then the countable dual group $M = \widehat{X}$ is a module over the ring R_d under the operation

$$f \cdot a = \sum_{\mathbf{n} \in \mathbb{Z}^d} f_{\mathbf{n}} \widehat{\alpha}(\mathbf{n})(a)$$

for $f \in R_d$ and $a \in M$. The module M is called the *dual module* of $\mathbf{X}$. Conversely, a countable module M over R_d determines an algebraic $\mathbb{Z}^d$ -action $\mathbf{X}_M = (X_M, \alpha_M)$ by setting

$$\widehat{\alpha}_M(\mathbf{n})(a) = \mathbf{u}^{\mathbf{n}} \cdot a$$

for every $\mathbf{n} \in \mathbb{Z}^d$ and $a \in M$.

An algebraic $\mathbb{Z}^d$ -action $\mathbf{X}$ is *Noetherian* if the dual module is Noetherian. The following properties are equivalent.

- $\mathbf{X}_M$ is Noetherian.
- M is finitely-generated over R_d (this is equivalent to M being Noetherian since R_d is itself Noetherian).
- Any descending chain of closed α_M -invariant subgroups of X_M stabilizes (the Descending Chain Condition; see [9]).

The topological entropy of the system $\mathbf{X}_M$ is defined and computed in terms of the module M in [11].

Rigidity properties of algebraic $\mathbb{Z}^d$ -actions have been studied by several authors. Measurable equivariant maps between mixing zero-entropy algebraic $\mathbb{Z}^d$ -actions exhibit strong regularity properties (see [3], [5], [8] and [10]). For a $\mathbb{Z}$ -action generated by an automorphism θ on a connected finite-dimensional compact abelian group, it is known that the topological centralizer of the action admits non-affine maps if and only if θ is not ergodic (cf. [1], [2] and [16]). Ergodic automorphisms of infinite-dimensional groups may have non-affine maps in their centralizers (see Example 1.3). In [4] it is shown that for any expansive connected algebraic $\mathbb{Z}^d$ -action $\mathbf{X}$, the topological centralizer of α consists of affine maps (expansiveness is a condition that implies the Descending Chain Condition; for $d = 1$ it forces the compact group X to be finite-dimensional).

In this note we prove the following result, which characterizes a form of topological rigidity in terms of topological entropy.

Theorem 1.1. *Let $\mathbf{X}_1, \mathbf{X}_2$ be connected mixing Noetherian algebraic $\mathbb{Z}^d$ -actions. Then the following properties are equivalent.*

- (1) *Every equivariant continuous map $\mathbf{X}_1 \rightarrow \mathbf{X}_2$ is an affine map.*
- (2) *The system $\mathbf{X}_2$ has finite topological entropy.*

For $d \geq 2$, this result applies to situations where the underlying group X_2 is infinite-dimensional, so the lifting techniques of [1] and [16]

cannot be applied directly. Write $\mathbb{T} \subset \mathbb{C}$ for the multiplicative unit circle.

Example 1.2. To illustrate Theorem 1.1, consider the $\mathbb{Z}^2$ -action $\mathbf{X}$ where $X \subset \mathbb{T}^{\mathbb{Z}^2}$ is the closed subgroup consisting of all $x \in \mathbb{T}^{\mathbb{Z}^2}$ with

$$x(m+1, n) \cdot x(m, n) \cdot x(m, n+1) = 1 \text{ for all } m, n \in \mathbb{Z},$$

and α is the shift action of $\mathbb{Z}^2$ on X . The system $\mathbf{X}$ is mixing and has finite entropy. It follows that every continuous equivariant map from $\mathbf{X}$ to itself is an affine map. In contrast, the *measurable* centraliser of $\mathbf{X}$ contains many non-affine maps, since $\mathbf{X}$ is measurably isomorphic to a $\mathbb{Z}^2$ Bernoulli shift (see [13], [17]).

Example 1.3. For the case of a single automorphism, the compact group being finite-dimensional forces the entropy to be finite. Ergodic automorphisms of infinite-dimensional groups are not topologically rigid in general.

For example, the shift automorphism of $X = \mathbb{T}^{\mathbb{Z}}$ defines an ergodic $\mathbb{Z}$ -action of infinite entropy that is not topologically rigid: if $f : \mathbb{T} \rightarrow \mathbb{T}$ is any map, then the shift map commutes with the map $\phi : X \rightarrow X$ defined by $(\phi(x))_k = f(x_k)$. The module corresponding to this action is a Noetherian R_1 -module.

On the other hand, an ergodic automorphism of $\mathbb{T}^{\mathbb{Z}}$ that splits into a direct product of automorphisms of finite-dimensional tori is topologically rigid. The module corresponding to this action is not Noetherian. It is not known whether such an action can have finite topological entropy (see [12]).

The next example again shows that Theorem 1.1 does not hold for non-Noetherian actions.

Example 1.4. Let F_d denote the field of fractions of R_d , considered as a R_d -module. Let $\mathbf{X}_1$ denote the algebraic $\mathbb{Z}^d$ -action corresponding to F_d . Notice that F_d is torsion-free as a R_d -module, and $\mathbf{X}_1$ has infinite entropy. For any $\mathbf{n} \in \mathbb{Z}^d$, multiplication by $\mathbf{u}^{\mathbf{n}} - 1$ is an automorphism of F_d . By duality, the map $x \mapsto \alpha_1(\mathbf{n})(x) - x$ is a continuous automorphism of X_1 for any $\mathbf{n} \in \mathbb{Z}^d$. In particular, $\mathbf{X}_1$ does not have any non-trivial periodic orbits. Now let $\mathbf{X}_2$ be any mixing connected algebraic $\mathbb{Z}^d$ -action with a dense set of periodic orbits (any Noetherian system has this property). Since continuous equivariant maps take periodic orbits to periodic orbits, it follows that any continuous equivariant map from $\mathbf{X}_2$ to $\mathbf{X}_1$ is trivial.

Example 1.4 is similar in spirit to a remark of Comfort (see [7]): there are no non-trivial homomorphisms $\mathbb{T} \rightarrow \widehat{\mathbb{Q}}$ since torsion elements are dense in $\mathbb{T}$ but absent in $\widehat{\mathbb{Q}}$.

2. ALGEBRAIC $\mathbb{Z}^d$ -ACTIONS

In this section basic results and terminology on algebraic $\mathbb{Z}^d$ -actions is collected. A prime ideal $\mathfrak{p} \subset R_d$ is *associated* with the R_d -module M if there exists $m \in M$ with $\mathfrak{p} = \{f \in R_d \mid f \cdot m = 0\}$. The set of prime ideals associated with M is denoted $\text{Asc}(M)$. If M is Noetherian, then $\text{Asc}(M)$ is finite. The torsion submodule of M is defined by

$$\text{Tor}(M) = \{m \in M \mid r \cdot m = 0 \text{ for some non-zero } r \in R_d\}.$$

A module M is said to be a *torsion module* if $\text{Tor}(M) = M$.

The following result taken from [14, Theorem 6.5] characterizes mixing in algebraic terms.

Lemma 2.1. *The algebraic $\mathbb{Z}^d$ -action X_M is mixing if and only if for every $\mathfrak{p} \in \text{Asc}(M)$ and for every non-zero $\mathbf{n} \in \mathbb{Z}^d$, the polynomial $\mathbf{u}^{\mathbf{n}} - 1$ does not lie in $\mathfrak{p}$.*

An algebraic $\mathbb{Z}^d$ -action X_2 is an *algebraic factor* of X_1 if there is a surjective continuous equivariant homomorphism $\phi : X_1 \rightarrow X_2$.

The next lemma shows that if the module corresponding to an algebraic $\mathbb{Z}^d$ -action is Noetherian, then infinite topological entropy can only be created by the presence of (a factor of) a full shift with infinite alphabet.

Lemma 2.2. *For a Noetherian system X_M the following conditions are equivalent.*

- (1) X_M does not admit a non-trivial closed α_M -invariant subgroup H with the property that the restriction of α_M to H is an algebraic factor of the shift action of $\mathbb{Z}^d$ on $(\mathbb{T}^n)^{\mathbb{Z}^d}$ for some $n > 0$.
- (2) M is a torsion module.
- (3) X_M has finite topological entropy.

Proof. (1) $\implies$ (2). Suppose that M is not a torsion module and $N = M / \text{Tor}(M)$. Then N is a non-zero torsion-free R_d -module.

We claim that N is isomorphic to a submodule of the free module R_d^n of rank n for some $n \geq 1$. Let N_0 denote the localisation of N at the prime ideal $\{0\}$. Since N is Noetherian, N_0 is a finite-dimensional vector space over $F = \mathbb{Z}(u_1^{\pm 1}, \dots, u_d^{\pm 1})$, the quotient field of R_d . Let $B = \{b_1, \dots, b_n\}$ be any F -basis of N_0 . The map $m \mapsto m/1$ embeds N as a submodule of N_0 . Choose a finite R_d -generating set A of N , and an element $p \in R_d$ with the property that $p \cdot a$ lies in the R_d -submodule generated by B for all $a \in A$. The submodule generated by $\{b_1/p, \dots, b_n/p\}$ contains N , and is a free R_d -module of rank n . This proves the claim.

The system X_N is therefore an algebraic factor of the shift action on $(\mathbb{T}^n)^{\mathbb{Z}^d}$. Since N is a quotient of M , by duality there exists a closed α -invariant subgroup $H \subset X$ such that the restriction of α to H is conjugate to X_N .

(2) $\implies$ (1). Let $H \subset X$ be a closed α -invariant subgroup such that the restriction of α to H is an algebraic factor of the shift action on $(\mathbb{T}^n)^{\mathbb{Z}^d}$ for some $n > 0$. The dual module of the shift action on $(\mathbb{T}^n)^{\mathbb{Z}^d}$ is isomorphic to the direct sum of n copies of R_d , which implies that $\widehat{H}$ is a torsion-free R_d -module. On the other hand, $\widehat{H}$ is a quotient of the R_d -module M , which is a torsion module by assumption (2). Hence $\widehat{H} = \{0\}$ so H is trivial.

(2) $\implies$ (3). Let $\{m_1, \dots, m_k\}$ generate M as an R_d -module. For $j = 1, \dots, k$ let $I_j \subset R_d$ denote the ideal defined by

$$I_j = \{p \in R_d \mid p \cdot m_j = 0\}.$$

Since M is a torsion module, each I_j is non-zero. For $j = 1, \dots, k$ let M_j denote the R_d -module R_d/I_j . Since each I_j is non-zero, $\mathbf{X}_{M_j}$ has finite entropy by [11, Theorem 3.1]. Let

$$M' = M_1 \oplus \dots \oplus M_k.$$

Since each $\mathbf{X}_{M_j}$ has finite entropy, $\mathbf{X}_{M'}$ also has finite entropy. The map $(r_1, \dots, r_k) \mapsto r_1 m_1 + \dots + r_k m_k$ expresses M as a quotient of M' . The dual of this map embeds $\mathbf{X}_M$ as a sub-action of $\mathbf{X}_{M'}$, so in particular $\mathbf{X}_M$ has finite entropy.

(3) $\implies$ (2). If M is not a torsion module, then it contains R_d as a submodule. By duality, the shift action of $\mathbb{Z}^d$ on $\mathbb{T}^{\mathbb{Z}^d}$ is therefore an algebraic factor of $\mathbf{X}_M$. Since the former action has infinite entropy, $\mathbf{X}_M$ has infinite topological entropy. $\square$

3. VAN KAMPEN'S THEOREM

In the proof of Theorem 1.1 the following structure theorem of van Kampen [15] will be used in place of the lifting of toral maps. This result splits continuous maps into a ‘linear’ part (a character) and a ‘non-linear’ part in a unique way. It is also used in this connection by Walters [16]. We include a short proof for the convenience of the reader.

Theorem 3.1. *Let X be a compact connected abelian group and let $f : X \rightarrow \mathbb{T}$ be a continuous map with $f(e) = 1$. Then there exist a character $\phi \in \widehat{X}$ and a continuous map $S(f) : X \rightarrow \mathbb{R}$ such that*

$$S(f)(e) = 0, \quad f(x) = \phi(x) \cdot e^{2\pi i S(f)(x)} \text{ for all } x \in X. \quad (1)$$

Moreover, ϕ and $S(f)$ are uniquely defined by (1).

Proof. Write $\exp(t) = e^{2\pi i t}$. We first prove that (1) determines ϕ and $S(f)$ uniquely. Suppose that

$$f = \phi_1 \cdot (\exp \circ h_1) = \phi_2 \cdot (\exp \circ h_2).$$

Then $\phi \cdot (\exp \circ h) = 1$, where $\phi = \phi_1 - \phi_2$ and $h = h_1 - h_2$. Define $p : X \times X \rightarrow \mathbb{R}$ by

$$p(x, y) = h(x + y) - h(x) - h(y).$$

Since ϕ and $\exp$ are homomorphisms and $\exp \circ h = \overline{\phi}$, it follows that $\exp \circ p$ is trivial, so the image of p is contained in $\mathbb{Z}$. Since X is connected, this implies that p is identically zero, so h is a continuous homomorphism from X to $\mathbb{R}$. This forces h to be identically zero (since $\mathbb{R}$ has no non-trivial compact subgroups), so $h_1 = h_2$ and $\phi_1 = \phi_2$.

Turning to the existence of ϕ and $S(f)$, we first consider two special cases.

- (1) If $X = \mathbb{T}^n$ for some n , then let $\pi_1(f) : \mathbb{Z}^n \rightarrow \mathbb{Z}$ be the induced map between fundamental groups. Choose $\phi \in \widehat{X}$ with the property that $\pi_1(f) = \pi_1(\phi)$. Since the map $f \cdot \overline{\phi}$ is null-homotopic and $\exp : \mathbb{R} \rightarrow \mathbb{T}$ is a covering map, there exists a unique continuous map $h : X \rightarrow \mathbb{R}$ such that $h(e) = 0$ and $f \cdot \overline{\phi} = \exp \circ h$.
- (2) If $f(X) \subset V = \exp(-1/4, 1/4)$ then, since the restriction of $\exp$ to $(-1/4, 1/4)$ is one-to-one, there exists a unique map

$$p : V \rightarrow (-1/4, 1/4)$$

such that $\exp \circ p$ is the identity on V . Thus $f = \exp \circ p \circ f$.

To prove the general case, choose a translation-invariant metric ρ on X . Since f is uniformly continuous, there exists $\delta > 0$ such that

$$\rho(x, y) < \delta \implies f(x)\overline{f(y)} \in V.$$

For any finite set $A \subset \widehat{X}$, let

$$X(A) = \{x \in X \mid \phi(x) = 1 \text{ for all } \phi \in A\}.$$

Then $X(A)$ is a closed subgroup of X since it is a finite intersection of kernels of characters. Moreover, the family of sets

$$\{X(A) \mid A \subset \widehat{X} \text{ is finite}\}$$

has the finite intersection property, and their intersection is $\{e\}$ since the character group separates points.

Let $Y \subset X$ denote the open subset defined by

$$Y = \{x \in X \mid \rho(e, x) < \delta/2\}.$$

Since $Y^c = X \setminus Y$ is compact, there exists a finite set $B \subset \widehat{X}$ such that $X(B) \subset Y$. For any $x \in X$, let

$$A_x = \{f(xz) \mid z \in X(B)\} \subset \mathbb{T}.$$

Since the diameter of A_x is strictly less than $1/2$, there exists a unique closed ball of minimal radius, $B_x \subset \mathbb{T}$, that contains A_x . Let $r(x)$ be the center of B_x . The map $x \mapsto r(x)$ is well-defined, continuous, and invariant under translations by elements of $X(B)$. Since X is compact

and connected, $\widehat{X}$ is torsion-free. On the other hand, the annihilator $X(B)^\perp$ is finitely-generated, so $X/X(B)$ is isomorphic to a torus. It follows that $r = \phi \cdot (\exp \circ p)$ for some $\phi \in \widehat{X}$ and some $p : X \rightarrow \mathbb{R}$ with $p(e) = 0$. Since $f \cdot \bar{r}(X) \subset V$, $f \cdot \bar{r} = \exp \circ q$ for some $q : X \rightarrow \mathbb{R}$ with $q(e) = 0$. Now $f = r \cdot (f \cdot \bar{r}) = \phi \cdot (\exp \circ (p + q))$. $\square$

4. RIGIDITY OF EQUIVARIANT MAPS

For any algebraic $\mathbb{Z}^d$ -action $\mathbf{X} = (X, \alpha)$ and for any locally compact abelian group A , denote by A^X the group of all continuous maps

$$h : X \rightarrow A, \quad h(e) = e,$$

equipped with point-wise multiplication. The action α induces the structure of an R_d -module on A^X by defining

$$p \cdot h(x) = \sum_{\mathbf{n} \in \mathbb{Z}^d} p(\mathbf{n}) \cdot h \circ \alpha(\mathbf{n})(x).$$

A key observation is that $\widehat{X}$ can be regarded as a submodule of $\mathbb{T}^X$ with this structure.

Proposition 4.1. *Let $\mathbf{X} = (X, \alpha)$ be a connected $\mathbb{Z}^d$ -action. Then $\mathbb{R}^X$ and $\mathbb{T}^X / \widehat{X}$ are isomorphic as R_d -modules.*

Proof. The correspondence $f \mapsto S(f)$ from Theorem 3.1 induces a map S from $\mathbb{T}^X$ to $\mathbb{R}^X$. If f_1, f_2 are elements of $\mathbb{T}^X$ then, by the uniqueness part of Theorem 3.1, $S(f_1 \bar{f}_2) = S(f_1) - S(f_2)$ so S is a group homomorphism. Similarly, if θ is a continuous endomorphism of G , then $S(f \circ \theta) = S(f) \circ \theta$. Hence $S : \mathbb{T}^X \rightarrow \mathbb{R}^X$ is an R_d -module homomorphism. For any f in $\mathbb{R}^X$, $S(e^{2\pi i f}) = f$, so the map S is surjective. Since $\ker(S) = \widehat{X}$, the statement follows. $\square$

If f and g are functions from $\mathbb{Z}^d$ to $\mathbb{C}$ and g has finite support, the convolution $f * g : \mathbb{Z}^d \rightarrow \mathbb{C}$ is given by

$$f * g(\mathbf{i}) = \sum_{\mathbf{j} \in \mathbb{Z}^d} f(\mathbf{i} - \mathbf{j}) \cdot g(\mathbf{j}).$$

Write $L^2(\mathbb{Z}^d)$ for the set of all square-integrable functions $\mathbb{Z}^d \rightarrow \mathbb{C}$ (with respect to the counting measure on $\mathbb{Z}^d$).

In addition to van Kampen's theorem, a simple version of the L^2 zero-divisor problem is needed (see [6] for an overview).

Proposition 4.2. *If $f \in L^2(\mathbb{Z}^d)$ has $f * g = 0$ for some non-zero function $g : \mathbb{Z}^d \rightarrow \mathbb{C}$ with finite support, then f is identically zero.*

Proof. Since the support of g is finite, there exists $\mathbf{n} \in \mathbb{Z}^d$ such that the support of $g * \delta_{\mathbf{n}}$ is contained in $\mathbb{N}^d$. Replacing g by $g * \delta_{\mathbf{n}}$ if necessary, we may assume that the support of g is contained in $\mathbb{N}^d$. Let $\widehat{f}, \widehat{g} \in L^2(\mathbb{T}^d)$ denote the Fourier transforms of f and g respectively. By the choice of

$g, \widehat{g} = p|_{\mathbb{T}^d}$ for some non-zero polynomial $p(\mathbf{z}) \in \mathbb{C}[z_1, \dots, z_d]$. Define $V(p) \subset \mathbb{T}^d$ by

$$V(p) = \{x \in \mathbb{T}^d \mid p(x) = 0\}.$$

We claim that $\lambda_d(V(p)) = 0$ for any non-zero p , where λ_d is Haar measure on $\mathbb{T}^d$. This may be proved by induction on d . If $d = 1$, then $V(p) \subset \mathbb{T}$ is finite since every non-zero polynomial has only finitely many roots. If $d > 1$, choose polynomials $p_0, \dots, p_k$ in $\mathbb{C}[z_1, \dots, z_{d-1}]$ such that

$$p(z_1, \dots, z_d) = \sum_{i=0}^k p_i(z_1, \dots, z_{d-1}) z_d^i.$$

Since p is non-zero, p_i is non-zero for some i . By the inductive hypothesis, $\lambda_{d-1}(V(p_i)) = 0$. If $(z_1, \dots, z_{d-1})$ lies in $\mathbb{T}^{d-1} \setminus V(p_i)$, then the map $z \mapsto p(z_1, \dots, z_{d-1}, z)$ is a non-zero polynomial in $\mathbb{C}[z]$. This implies that for any $(z_1, \dots, z_{d-1}) \in \mathbb{T}^{d-1} \setminus V(p_i)$, the set

$$\{z \in \mathbb{T} \mid (z_1, \dots, z_{d-1}, z) \in V(p)\}$$

is finite. By Fubini's theorem,

$$\lambda_d(V(p)) = \int_{\mathbb{T}^{d-1}} \int_{\mathbb{T}} \mathbb{I}_{V(p)} d\lambda_1 d\lambda_{d-1} = 0,$$

where $\mathbb{I}$ is the indicator function, which proves the claim. Since $\widehat{g} = p|_{\mathbb{T}^d}$ and $\widehat{f} \cdot \widehat{g} = \widehat{f * g} = 0$, it follows that $\widehat{f} = 0$ almost everywhere, so $f = 0$. $\square$

Lemma 4.3. *If $\mathbf{X} = (X, \alpha)$ is a mixing connected algebraic $\mathbb{Z}^d$ -action, then $\text{Tor}(\mathbb{T}^X) \subset \widehat{X}$.*

Proof. Let M denote the set of all square-integrable functions

$$h : \widehat{X} \rightarrow \mathbb{C}, \quad h(e) = 0.$$

Defining

$$p \cdot h(\phi) = \sum_{\mathbf{i} \in \mathbb{Z}^d} p(\mathbf{i}) h(\phi \circ \alpha(\mathbf{i}))$$

for $p \in R_d$ gives M the structure of an R_d -module.

We claim first that M is torsion-free. Let h be an element of $\text{Tor}(M)$; for any non-trivial $\chi \in \widehat{X}$ define a function $h_\chi : \mathbb{Z}^d \rightarrow \mathbb{C}$ by

$$h_\chi(\mathbf{i}) = h(\chi \circ \alpha(\mathbf{i})).$$

Since α is mixing, the map $\mathbf{i} \mapsto \chi \circ \alpha(\mathbf{i})$ is one-to-one. Hence

$$\sum_{\mathbf{i} \in \mathbb{Z}^d} |h_\chi(\mathbf{i})|^2 \leq \sum_{\chi \in \widehat{X}} |h(\chi)|^2 < \infty.$$

This shows that $h_\chi \in L^2(\mathbb{Z}^d)$ for all $\chi \in \widehat{X}$. Note that $L^2(\mathbb{Z}^d)$ itself is an R_d -module with respect to the multiplication $p \cdot h = p * h$. Furthermore, the map $h \mapsto h_\chi$ is an R_d -module homomorphism from M to $L^2(\mathbb{Z}^d)$.

Since $\text{Tor}(L^2(\mathbb{Z}^d)) = \{0\}$ by Proposition 4.2, we conclude that $h_\chi = 0$ for all χ , so $h = 0$. This proves that M is torsion-free.

For f in $\mathbb{R}^X$, let $\widehat{f} \in M$ denote the Fourier transform of f . Since $\widehat{f \circ \theta}(\phi) = \widehat{f}(\phi \circ \theta)$ for any continuous endomorphism θ of X , the map $f \mapsto \widehat{f}$ is an R_d -module homomorphism from $\mathbb{R}^X$ to M . By the Fourier inversion theorem this map is injective. Since $\text{Tor}(M) = \{0\}$, this implies that $\text{Tor}(\mathbb{R}^X) = \{0\}$. Proposition 4.1 then shows that $\text{Tor}(\mathbb{T}^X) \subset \widehat{X}$. $\square$

We are now ready to prove Theorem 1.1.

Proof. Suppose that X_2 has finite entropy, and let f be an equivariant continuous map $X_1 \rightarrow X_2$. Define $f_0 : X_1 \rightarrow X_2$ by

$$f_0(x) = f(x) - f(e). \quad (2)$$

Since f is equivariant, so is f_0 .

Fix an arbitrary character $\phi \in \widehat{X_2}$. By Lemma 2.2, $\widehat{X_2}$ is a torsion module, so ϕ lies in the torsion submodule of $\mathbb{T}^{X_2}$. Since f_0 is equivariant and $f_0(e) = 1$, the map $h \mapsto h \circ f_0$ is an R_d -module homomorphism $\mathbb{T}^{X_2} \rightarrow \mathbb{T}^{X_1}$. Hence $\phi \circ f_0$ is an element of the torsion submodule of $\mathbb{T}^{X_1}$. By Lemma 4.3, $\phi \circ f_0$ lies in $\widehat{X_1}$. Since the initial choice of ϕ was arbitrary, this shows that $\phi \mapsto \phi \circ f_0$ is a group homomorphism from $\widehat{X_2}$ to $\widehat{X_1}$. By duality, there exists a continuous homomorphism $\theta : X_1 \rightarrow X_2$ such that $\phi \circ f_0 = \phi \circ \theta$ for all $\phi \in \widehat{X_2}$. Since characters separate points, this implies that $f_0 = \theta$. Hence $f = f(e) + f_0$ is an affine map.

If X_2 has infinite entropy, then by Lemma 2.2 there exists a non-trivial closed α_2 -invariant subgroup $H \subset X_2$ with the property that the restriction of α_2 to H is an algebraic factor of the shift action on $(\mathbb{T}^n)^{\mathbb{Z}^d}$ for some $n > 0$. Let $K \subset (\mathbb{T}^n)^{\mathbb{Z}^d}$ be a proper, closed shift-invariant subgroup such that the restriction of α_2 to H is algebraically conjugate to the shift action of $\mathbb{Z}^d$ on $(\mathbb{T}^n)^{\mathbb{Z}^d}/K$. Since any non-affine equivariant map from X_1 to H gives rise to a non-affine equivariant map $X_1 \rightarrow X_2$, without loss of generality we may assume that $X_2 = (\mathbb{T}^n)^{\mathbb{Z}^d}/K$, and α_2 is the shift action.

For any continuous map $q : X_1 \rightarrow \mathbb{R}^n$ define a map

$$\sigma(q) : X_1 \rightarrow (\mathbb{T}^n)^{\mathbb{Z}^d}$$

by

$$\sigma(q)(x)(\mathbf{n}) = \exp \circ q \circ \alpha_1(\mathbf{n})(x).$$

Let $\pi : (\mathbb{T}^n)^{\mathbb{Z}^d} \rightarrow (\mathbb{T}^n)^{\mathbb{Z}^d}/K$ denote the projection map. For any

$$q : X_1 \rightarrow \mathbb{R}^n,$$

$\pi \circ \sigma(q)$ is a continuous equivariant map from X_1 to X_2 . We claim that there exists a non-zero continuous equivariant map from X_1 to X_2 of the form $\pi \circ \sigma(q)$ for some continuous map $q : X_1 \rightarrow \mathbb{R}^n$ with $q(e) = 0$.

For any finite set $F \subset \mathbb{Z}^d$, let Π_F denote the projection map

$$\Pi_F : (\mathbb{T}^n)^{\mathbb{Z}^d} \rightarrow (\mathbb{T}^n)^F.$$

Since K is a proper closed subgroup of $(\mathbb{T}^n)^{\mathbb{Z}^d}$, there exists a finite set $F \subset \mathbb{Z}^d$, and a point $\mathbf{x} \in (\mathbb{T}^n)^F$, such that $\mathbf{x}$ does not lie in the image of Π_F . Since X_1 is mixing, for any $\mathbf{i} \neq \mathbf{j} \in \mathbb{Z}^d$, the kernel of $\alpha_1(\mathbf{i}) - \alpha_1(\mathbf{j})$ is a proper closed subgroup of X_1 . In particular, there exists $y \in X_1$ such that $y \neq e$, and

$$\alpha_1(\mathbf{i})(y) \neq \alpha_1(\mathbf{j})(y) \text{ for any } \mathbf{i}, \mathbf{j} \in F.$$

Choose $\mathbf{z} \in (\mathbb{R}^n)^F$ such that $\exp(\mathbf{z}(\mathbf{i})) = \mathbf{x}(\mathbf{i})$ for all $\mathbf{i} \in F$. Let

$$q : X_1 \rightarrow \mathbb{R}^n$$

be any continuous map with $q(e) = 0$ and $q \circ \alpha_1(\mathbf{i})(y) = \mathbf{z}(\mathbf{i})$ for all $\mathbf{i} \in F$. Since $\pi \circ \sigma(q)(y)$ does not lie in K , this proves the claim.

Now let $q : X_1 \rightarrow \mathbb{R}^n$ be any continuous map such that $q(e) = 0$, and $\pi \circ \sigma(q) : X_1 \rightarrow X_2$ is a non-zero map. For any $t \in [0, 1]$, define maps $q_t : X_1 \rightarrow \mathbb{R}^n$ and $h_t : X_1 \rightarrow X_2$ by $q_t(x) = tq(x)$, $h_t(x) = \pi \circ \sigma(q_t)$. For any $t \in [0, 1]$, h_t is a continuous equivariant map from X_1 to X_2 , and $h_t(e) = e$. We claim that h_t is non-affine for some $t \in (0, 1]$.

Suppose this is not the case. Then for each $t \in [0, 1]$, h_t is a continuous homomorphism from X_1 to X_2 . Let Y denote the set of all continuous maps from X_1 to X_2 . Choose any metric ρ on X_2 that gives the topology, and define a metric ρ_0 on Y by

$$\rho_0(h_1, h_2) = \sup\{\rho(h_1(x), h_2(x)) \mid x \in X_1\}.$$

The map $t \mapsto h_t$ is continuous with respect to ρ_0 and the set of all continuous homomorphisms from X_1 to X_2 forms a discrete subset of Y . Hence $t \mapsto h_t$ is constant, which contradicts the fact that $h_0 = 0$ and $h_1 \neq 0$. This proves that some h_t is not an affine map. Since h_t is a continuous equivariant map from X_1 to X_2 for any $t \in [0, 1]$, Theorem 1.1 follows. $\square$

REFERENCES

1. R. L. Adler and R. Palais, *Homeomorphic conjugacy of automorphisms on the torus*, Proc. Amer. Math. Soc. **16** (1965), 1222–1225.
2. D. Z. Arov, *Topological similitude of automorphisms and translations of compact commutative groups*, Uspehi Mat. Nauk **18** (1963), no. 5 (113), 133–138.
3. S. Bhattacharya, *Higher order mixing and rigidity of algebraic actions on compact abelian groups*, Israel J. Math. (to appear), www.esi.ac.at/Preprint-shadows/esi1128.html.
4. ———, *Orbit equivalence and topological conjugacy of affine actions on compact abelian groups*, Monatsh. Math. **129** (2000), no. 2, 89–96.

5. S. Bhattacharya and K. Schmidt, *Homoclinic points and isomorphism rigidity of algebraic $\mathbb{Z}^d$ -actions on zero-dimensional compact abelian groups*, Israel J. Math. (to appear), www.esi.ac.at/Preprint-shadows/esi1127.html.
6. D. R. Farkas and P. A. Linnell, *Zero divisors in group rings: something old, something new*, Representation theory, group rings, and coding theory, Contemp. Math., vol. 93, Amer. Math. Soc., Providence, RI, 1989, pp. 155–166.
7. G. L. Itzkowitz, *The existence of homomorphisms in compact connected Abelian groups*, Proc. Amer. Math. Soc. **19** (1968), 214–216.
8. A. Katok, S. Katok, and K. Schmidt, *Rigidity of measurable structure for $\mathbb{Z}^d$ -actions by automorphisms of a torus*, Comment. Math. Helv. (to appear), www.esi.ac.at/Preprint-shadows/esi850.html.
9. B. Kitchens and K. Schmidt, *Automorphisms of compact groups*, Ergodic Theory Dynam. Systems **9** (1989), no. 4, 691–735.
10. ———, *Isomorphism rigidity of irreducible algebraic $\mathbb{Z}^d$ -actions*, Invent. Math. **142** (2000), no. 3, 559–577.
11. D. Lind, K. Schmidt, and T. Ward, *Mahler measure and entropy for commuting automorphisms of compact groups*, Invent. Math. **101** (1990), no. 3, 593–629.
12. D. A. Lind, *Ergodic automorphisms of the infinite torus are Bernoulli*, Israel J. Math. **17** (1974), 162–168.
13. D. J. Rudolph and K. Schmidt, *Almost block independence and Bernoullicity of $\mathbb{Z}^d$ -actions by automorphisms of compact abelian groups*, Invent. Math. **120** (1995), no. 3, 455–488.
14. K. Schmidt, *Dynamical systems of algebraic origin*, Progress in Mathematics, vol. 128, Birkhäuser Verlag, Basel, 1995.
15. E. R. van Kampen, *On almost periodic functions of constant absolute value*, J. London Math. Soc. **12** (1937), 3–6.
16. P. Walters, *Topological conjugacy of affine transformations of compact abelian groups*, Trans. Amer. Math. Soc. **140** (1969), 95–107.
17. T. Ward, *Almost block independence for the three dot $\mathbb{Z}^2$ dynamical system*, Israel J. Math. **76** (1991), no. 1-2, 237–256.

SB: SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, BOMBAY 400005, INDIA

E-mail address: `siddhart@math.tifr.res.in`

TW: SCHOOL OF MATHEMATICS, UNIVERSITY OF EAST ANGLIA, NORWICH NR4 7TJ, ENGLAND

E-mail address: `t.ward@uea.ac.uk`